



**ENGLISH TITLE (USE A CAPITALIZE 14 PT BOLD,  
AND A TRIANGULAR HEAD IF MORE THAN  
ONE LINE FORMAT)**



**FONT SIZE  
12 pt BOLD**

**NAME LASTNAME**

Spacing balance

\* MASTER/ DOCTOR

**MASTER OF NAME**

Spacing balance

**IN**

**PROGRAM NAME**

Spacing balance


**SCHOOL OF NAME**

**MAE FAH LUANG UNIVERSITY**



**Academic year of submission**

**©COPYRIGHT BY MAE FAH LUANG UNIVERSITY**

**Font size  
10 pt BOLD**



**ENGLISH TITLE SAME COVER**



1.5 inch



1 inch



**NAME LASTNAME**



THESIS  
DISSERTATION  
INDEPENDENT STUDY



**THIS THESIS IS A PARTIAL FULFILLMENT OF  
THE REQUIREMENTS FOR THE DEGREE OF**

**MASTER OF NAME**

\* MASTER/ DOCTOR



**IN**

**PROGRAM NAME**

**FONT SIZE  
12 pt BOLD**



**SCHOOL OF NAME**


**MAE FAH LUANG UNIVERSITY**


**Academic year of submission**

**©COPYRIGHT BY MAE FAH LUANG UNIVERSITY**



**Font size  
10 pt BOLD**





2 inches

Font Size 14 Bold

## ACKNOWLEDGEMENTS

Font size 12

Spacing 24 pt

0.5 inches Writing an acknowledgement is the last and the best part of the dissertation writing process. It is a statement written by the author acknowledging the persons who assisted in writing the dissertation by being involved in different ways. You have the choice in your dissertation to leave out the acknowledgement, but this hardly ever happens. There has to be a way for you to thank the persons who helped you, made sacrifices in terms of time and effort and supported you to complete your dissertation successfully.

1.5 inches

1 inch

Spacing 12 pt

After all it is not the proper etiquette as not to do so. Acknowledgments do not need to comply with any fixed format or structure. It should be written in simple and in a personal tone. But sometimes, simpler matters turn out to be more difficult than one assumes.

Spacing 24 pt

Name Lastname

(3)

1 inch

THESIS  
DISSERTATION  
INDEPENDENT STUDY

2 inches

Title, use a capitalize only the first letter of word, except proper nouns.

**Thesis Title**

English Title

Spacing 12 pt

**Author**

Name Lastname

Spacing 12 pt

**Degree**

Master of ..... (Name of Program)

Spacing 12 pt

**Advisor**

Academic Position Name Lastname, Degree

**Co-Advisor (if any)**

Academic Position Name Lastname, Degree

Academic Position Name Lastname, Degree

Spacing 24 pt

**ABSTRACT**

Font size 14/ Bold

-0.5 inch-

Describing the abstract as a summary is accurate, but is also an understatement. Generally, an abstract contains the thesis of research conducted, the methods through which the thesis was tested, the findings of the research, and any final thoughts. An abstract should be as succinct as possible, and readers should not have difficulty comprehending the purpose of the research.

1.5 inches

1 inch

Spacing 12 pt

According to the APA, abstract length is limited to 120 words. Abstracts longer than 120 words may be truncated when included in databases, limiting searchability. The information included in an abstract depends on the type of paper. A report of an empirical study should contain a statement of the problem, subjects, method, findings and conclusion.

Spacing 24 pt

Font size 12/ Bold

**Keywords:** Keyword, Keyword, Keyword, Keyword, Keyword, Keyword, Keyword, Keyword, Keyword, Keyword, Keyword, Keyword

Font size 12 pt

(4)

1 inch

2 inches

Font size 14 / Bold

# TABLE OF CONTENTS

Spacing 24 pt

**Page**

Spacing 12 pt

**ACKNOWLEDGEMENTS**

**(3)**

**ABSTRACT**

**(4)**

**LIST OF TABLES**

**(Page)**

**LIST OF FIGURES**

**(Page)**

Spacing 12 pt

**CHAPTER**

**1 INTRODUCTION**

**1**

1.1 Title

Page

1.2 Title

Page

1.3 Title

Page

1.4 Title

Page

1.5 Title

Page

1.No. Title

Page

Spacing 12 pt

**2 Title**

**Page**

2.1 Title

Page

2.2 Title

Page

2.3 Title

Page

2.4 Title

Page

Spacing 12 pt

**3 Title**

**Page**

3.1 Title

Page

3.2 Title

Page

(Page No.)

1 inch

## TABLE OF CONTENTS (continued)

	<b>Page</b>
<b>CHAPTER</b>	
3.3 Title	Page
3.No. Title	Page
<b>4 Title</b>	Page
4.1 Title	Page
4.2 Title	Page
4.3 Title	Page
4.No. Title	Page
<b>5 Title</b>	<b>Page</b>
5.1 Title	Page
5.2 Title	Page
5.3 Title	Page
5.No. Title	Page
<b>REFERENCES</b>	<b>Page</b>
<b>APPENDICES</b>	<b>Page</b>
APPENDIX A NAME	Page
APPENDIX B NAME	Page
APPENDIX C NAME	Page
<b>CURRICULUM VITAE</b>	<b>Page</b>

## LIST OF TABLES

<b>Table</b>	<b>Page</b>
1.1 Frame Work	Page
No. Title	Page
No. Title	Page
No. Title	Page
No. Title	Page
No. Title	Page
No. Title	Page
No. Title	Page
No. Title	Page
No. Title	Page
No. Title	Page
No. Title	Page
No. Title	Page
No. Title	Page
No. Title	Page
No. Title	Page
No. Title	Page
No. Title	Page
No. Title	Page
No. Title	Page
No. Title	Page



## LIST OF FIGURES

<b>Figure</b>	<b>Page</b>
2.1 Type of Skin	Page
No. Title	Page
No. Title	Page
No. Title	Page
No. Title	Page
No. Title	Page
No. Title	Page
No. Title	Page
No. Title	Page
No. Title	Page
No. Title	Page
No. Title	Page
No. Title	Page
No. Title	Page
No. Title	Page
No. Title	Page
No. Title	Page
No. Title	Page
No. Title	Page
No. Title	Page
No. Title	Page
No. Title	Page

## **ABBREVIATIONS AND SYMBOLS**

VT	Vermont
VA	Virginia
WA	Washington
WV	West Virginia
WI	Wisconsin
WY	Wyoming

2 inch

## CHAPTER 1

Spacing 24 pt

### INTRODUCTION

Spacing 24 pt

Topic : use Capitalize Each Word

#### 1.1 Introduction to the Research Problem and Related Technologies in Fresh-Cut Mango

Spacing 24 pt

**-0.5 inch** → The global production of major tropical fruits such as mango, mangosteen, guava, pineapple, avocado and papaya was estimated at about 100 million tonnes in 2018. The world production of mango accounted for 52% of total global major tropical fruit production (Altendorf, 2019). Mango (*Mangifera indica* L.) is one of the most consumed tropical fresh fruits globally. Asia is the largest mango producing zone, especially India, China, Thailand, Pakistan, Indonesia, Bangladesh and the Philippines are world-leading mango production countries (Evans et al., 2017).

1.5 inch

Thailand has more than 60 mango varieties but the major varieties such as ‘Nam Dok Mai’, ‘Khiao Sawoei’, ‘Chok Anan’, ‘Nang Klang Wan’, ‘Pim Sen Dang’, ‘Maha Chanok’ and ‘Rad’ are popular in foreign markets. ‘Nam Dok Mai’ mango is the most popular variety for export to Japan and the global markets (Yasunaga et al., 2018). ‘Nam Dok Mai’ mango is one of Thailand’s most significant mangoes for both local and export markets due to its characteristics of great flavor and bright yellow flesh (Matulaprungsan et al., 2019). Mango possesses a unique flavor, aroma, texture and color and a high amount of bioactive compounds that attract consumers (Salinas-Roca et al., 2017).

1 inch

**-0.5 inch** → The variation in pre- and postharvest factors affect the quality and contribute to providing potent heterogeneous batches of mangoes in terms of fruit size, gustatory quality, essential nutrients, vitamins, and minerals, in the supply chain and affect the postharvest management (Jacobi et al., 1995; Lalel et al., 2003; Nunes et al., 2007). A study on postharvest losses of mango at different stages from harvesting to consumption revealed that these losses accounted for about 34.49 % in India.

1 inch



Sab et al. (2017) reported that the losses were 8.44 % at farm level, 4.93 % at the wholesale market level, 5.46 % at the retail market level, 5.65 % at the storage level, 3.19 % at processing and 6.82 % at the consumer level. Yusuf et al. (2019) also reported that the overall postharvest losses of mango from harvesting to consumption were accounted for about 25.51 % in Bangladesh, including 7.04 % losses at farm level, 4.7 % at wholesale market, 3.66 % at retail market, 3.5 % at storage, 3.5 % at consumer level, and 3.11 % at processing. Over-ripening, withering, and spoilage are the main postharvest losses at the retail and consumer level.

1 inch

1.5 inch

### References

If use **Name-Year citation**  
Example: Sort A to Z in references list  
Follow APA style

Altendorf, S. (2019). *Major tropical fruits market review 2018*. FAO.

Evans, E. A., Ballen, F. H., & Siddiq, M. (2017). Mango production, global trade, consumption trends, and postharvest processing and nutrition. In M. Siddiq, J. K. Brecht, & J. S. Sidhu (Eds.), *Handbook of Mango Fruit* (pp. 1-16). John Wiley & Sons.

### References


If use **Numerical citation**  
Example: Sort [1] to [last number] in references list  
Follow APA style


[1] Altendorf, S. (2019). *Major tropical fruits market review 2018*. FAO.

[2] Evans, E. A., Ballen, F. H., & Siddiq, M. (2017). Mango production, global trade, consumption trends, and postharvest processing and nutrition. In M. Siddiq, J. K. Brecht, & J. S. Sidhu (Eds.), *Handbook of Mango Fruit* (pp. 1-16). John Wiley & Sons.


If want to make a reference list at the end of the last chapter.

This can be done by making it like a normal thesis format. [Click here](#)







## CHAPTER 2




### LITERATURE REVIEW




#### 2.1 Mango




 Mango (*Mangifera indica* L.) belongs to the family *Anacardiaceae*, and it is originated in the Indo-Burma region. *Mangifera indica* can be distinguished into two groups, a subtropical group with mono-embryonic seed (Indian type) and a tropical group with poly-embryonic seed, depending on their manner of reproduction as well as their specific diversity areas (South-east Asian) (Litz, 2008). Mangoes are cultivated in tropical and subtropical locations under a variety of environmental conditions. Asia is the biggest mango production zone, and production was about 77.17 % of total world's mango production (Evans et al., 2017; FAOSTAT, 2015). It is one of the most important tropical fruit crops. More than a thousand mango varieties can be classified depending on shapes, sizes, colors, textures, and nutritional properties (Evans et al., 2017).

1.5 inch

1 inch


 The mango fruits possess a massive, fleshy drupe with a varied thickness edible mesocarp. The mesocarp is resinous and has a wide range of structure, size, color, fiber content, and flavor. Fruits vary in shape, forming elongate, oblong, and ovate or intermediate shapes combining two of these types. The length of fruits ranges from 2.5 to greater than 30 cm, depending on the varieties (Mukherjee & Litz, 2009). The most popular varieties of mango traded are Tommy Atkins, Kent, Keitt, Haden, Francis, Ataulfo, Alphonso, Kesar, Edward and Manila. Other varieties gaining importance in the trade include Sindhri, Badami, Glenn, Valencia Pride and Nam Dok Mai (Evans et al., 2017). In Thailand, one of the well-known kinds of mango is 'Nam Dok Mai Si-Thong (golden variety) which is highly produced in Thailand for domestic and export (Figure 2.1). The mangoes are majorly produced in Chiang Mai, Phitsanulok, Loei, Nakhon Ratchasima and Prachuap Khiri Khan provinces. The on-season production befalls from April to May, early off-season production is from January to March, and late off-season is from August to December (Statistics, 2015).



## 2.2 Problems of Fresh-Cut Mango Production

→ Fruit and vegetable processing is readily perishable and has a reduced shelf life than whole fruit because peeling, trimming, chopping, and shredding put them under a lot of physical stress (Kim, 2008). Many physiological changes can occur during the processing, including increased respiration rate, oxidation process, changing in color, and loss of nutrients because of several biochemical and enzymatic processes. Several enzymatic processes induce color and texture deterioration (Kong & Singh, 2016). The raw material quality is closely related to fresh-cut quality, so the raw mango attributes must be monitored before processing (Siddiq et al., 2017).

→ Physiological changes in wounded tissues tend to degrade cell integrity and subcellular compartmentation during fresh-cut processing, which contributes to endogenous enzyme and its substrates interact in a variety of ways. Wounding accelerates cell wall enzyme activity (Karakurt & Huber, 2002), inducing a great of tissues softening in fresh-cuts. These changes are responses to plant tissue injury during processing. Certain reactions to wounding include changes in chemical or physical properties, such as increased ion leakage or partial loss, turgor reduction and flux subsequent changes. These physiological changes bring declining texture and undesired essence, fresh-cut surface mottling, subsequent nutrient reduction, and excessive leaching moisture at the cut surface, that speeds up water depletion (Temiz & Kütük Ayhan, 2017).



Note : Close Figure/Table →

Note {Non-wounded —, wounded - - - }

Source Saltveit (2020)

**Figure 2.1** Effect of wounding on the respiration and ethylene production rate of climacteric fruits

The second line corresponds to the name on the first line.

**References**

If use **Name-Year citation**  
Example: Sort A to Z in references list  
Follow APA style

1.5 inch

1.5 inch

1 inch

Altendorf, S. (2019). *Major tropical fruits market review 2018*. FAO.

Litz, R. E. (2008). Mango. In C. Kole & T. C. Hall (Eds.), *Compendium of transgenic crop plants: Transgenic tropical and subtropical fruits and nuts* (pp. 163-174). Blackwell.

**References**

If use **Numerical citation**  
Example: Sort [1] to [last number] in references list  
Follow APA style

[1] Litz, R. E. (2008). Mango. In C. Kole & T. C. Hall (Eds.), *Compendium of transgenic crop plants: Transgenic tropical and subtropical fruits and nuts* (pp. 163-174). Blackwell.

[2] Altendorf, S. (2019). *Major tropical fruits market review 2018*. FAO.

If want to make a reference list at the end of the last chapter.

This can be done by making it like a normal thesis format. [Click here](#)

1 inch

2 inch

1 inch

### CHAPTER 3

Spacing 24 pt

## INVESTIGATION AND EVALUATION OF IMPACT BRUISING IN GUAVA USING IMAGE PROCESSING AND RESPONSE

1.5 inch

### SURFACE METHODOLOGY<sup>1</sup>

Spacing 24 pt

<sup>1</sup> = Enter a number to indicate the footnote

#### Abstract

-0.5 inch-

→ Simulated impact damage testing was investigated by fractal image analysis using response surface methodology (RSM) with a central composite design (CCF) on quality of ‘Glom Sali’ guava for drop heights (0.2, 0.4, and 0.6 m), number of drops (1, 3, and 5) and storage temperature conditions (10, 20, and 30 °C). After 48 h, impacted fruit were determined and analyzed for bruise area (BA), bruise volume (BV), browning index (BI), total color difference ( $\Delta E$ ), image analysis for bruise area (BAI), and fractal dimension (FD) at the bruising region on peeled guava. Results showed that the correlation coefficient ( $r = -0.6055$ ) between  $\Delta E$  and FD value was higher than  $\Delta E$  and either BA ( $r = 0.3132$ ) or BV ( $r = 0.2095$ ). The FD variable was determined as a better indicator than conventional measurement (BA or BV) for pulp browning and impact bruising susceptibility. The FD variable also exhibited highest  $R^2_{adj}$  value (81.69%) among the other five variables, as the highest precision model with high determination coefficient value ( $R^2_{adj}$ ) ( $> 0.8$ ) for impact bruising prediction. Recommended condition of the FD variable to minimize impact bruising was drop height of 0.53 m for five drops under storage at 30 °C. FD variable assessed by image analysis was shown to be a highly capable measurement to determine impact bruising susceptibility in guava fruit.

**Keywords:** Bruise Susceptibility, Impact Bruise, Mechanical Injury, Transportation

Add Line 2 inch

-0.8 inch-

→ <sup>1</sup>This paper has been published in *Horticulturae Journal*, 2021, 7(10), 411. <https://doi.org/10.3390/horticulturae7100411>



### 3.1 Introduction

Guava (*Psidium guajava* L.) is one of the most famous and economically important fruits in Thailand, with an export value in 2020 worth 5.50 million USD [1]. Guava is a climacteric fruit with a round shape and thin skin that bruises easily. To maintain fruit quality and shelf life and minimize losses, guava fruit needs proper postharvest handling practices [2]. Thai agriculture and ASEAN standards of guava require slight defects on the skin not exceeding 10% of the total surface area of guava fruit [3,4]. Bruising effects can be distinguished from quality changes in guava such as browning, softening of the fruit peel, cell destruction, and reduction in intercellular air spaces resulting in the bruised tissue losing moisture and becoming desiccated [5]. Impact damage to fruit is more severe than vibration and compression damages. When a fruit falls with sufficient force against a surface, impact damage occurs, while dynamic damage of a single fruit occurs through fruit-to-fruit impact between packaging. Fruit dropping from trees to the ground during harvesting, dynamic impact between single fruit, and between the fruit and packaging or containers are all causes of impact damage [6].

#### 3.1.1 Quality Measurements

Text.....

##### 2.2.2.1 Bruise Determination of the Guava Fruit

##### 2.2.2.2 Pulp Color at Bruise Area

1. Text....

2. Text....

1) Text.....

A. Text....

B. Text....

a. Text.....

**Table 3.1** Bruise Assessment Parameters of Guava Fruit Stored at 25 °C Under 70% RH for Four Days


Spacing 24 pt

	<b>BA</b>	<b>BV</b>	<b>BI</b>	$\Delta E$	<b>BAI</b>	<b>FD</b>
BA	1.0000					
BV	0.2363					
BI	-0.1871	0.1332				
$\Delta E$	0.3132	0.2095	0.4332			
BAI	0.9975 *	0.2372	-0.1837	0.3116		
FD	-0.3854	-0.2366	-0.0448	-0.6055 *	-0.3992	1.0000

Head Bold

**Note** \* Significant at  $p < 0.05$ : BA = bruise area; BV = bruise volume; BI = browning index;  $\Delta E$  = total color difference, BAI = bruise area by image analysis; FD = fractal dimension.

Note: No space / close the table  
If have source add under note 24 pt



**Figure 3.1** Verification of Predictive Model Equations for Fractal Dimension (FD)

Values in 'Glom Sali' Bruised Guava by Impact Testing for 48 h

If you have note/source in figure order  
Note  
Source

Spacing 24 pt

1 inch

## Acknowledgments

The authors gratefully acknowledge the Thailand International Cooperation Agency (TICA) and the Integrated AgriTech Ecosystem Research Group (IATE), Scientific and Technological Instruments Center, Mae Fah Luang University for financial and equipment support.

## References

If use **Numerical citation**

Example: Sort [1] to [last number] in references list

Follow APA style

[1] Thai Customs. (2021). *The exported guava from Thailand from 2015–2019*.

[http://www.customs.go.th/statistic\\_report.php?lang=en&ini\\_content=statistics\\_report&fbclid=IwAR3fgvhbyliB4O9LUuMedqNtqYNS\\_ZfyBOOqjPJnBGQlXT6zEWd18v577YQ](http://www.customs.go.th/statistic_report.php?lang=en&ini_content=statistics_report&fbclid=IwAR3fgvhbyliB4O9LUuMedqNtqYNS_ZfyBOOqjPJnBGQlXT6zEWd18v577YQ)

[2] Shafie, M., Rajabipour, A., & Mobli, H. (2017). Determination of bruise incidence of pomegranate fruit under drop case. *International Journal of Fruit Sciences*, *17*, 1–14.

[3] Li, Z., & Thomas, C. (2014). Quantitative evaluation of mechanical damage to fresh fruits. *Trends Food Science & Technology*, *35*, 138–150.

## References

If use **Name-Year citation**

Example: Sort A to Z in references list

Follow APA style

Li, Z., & Thomas, C. (2014). Quantitative evaluation of mechanical damage to fresh fruits. *Trends Food Science & Technology*, *35*, 138–150.

Shafie, M., Rajabipour, A., & Mobli, H. (2017). Determination of bruise incidence of pomegranate fruit under drop case. *International Journal of Fruit Sciences*, *17*, 1–14.

Thai Customs. (2021). *The exported guava from Thailand from 2015–2019*.

[http://www.customs.go.th/statistic\\_report.php?lang=en&ini\\_content=statistics\\_report&fbclid=IwAR3fgvhbyliB4O9LUuMedqNtqYNS\\_ZfyBOOqjPJnBGQlXT6zEWd18v577YQ](http://www.customs.go.th/statistic_report.php?lang=en&ini_content=statistics_report&fbclid=IwAR3fgvhbyliB4O9LUuMedqNtqYNS_ZfyBOOqjPJnBGQlXT6zEWd18v577YQ)


**APPENDIX**

for one appendix



or

**APPENDICES**

for many appendices



\* If you have only one appendix, do not type A



**APPENDIX B**



**CLINICAL ASSESSMENT OF  
REPIGMENTATION FORM**

Assessment of Repigmentation (CAR)

Please indicate your choice by applying (X);

<b>Patient no. 1</b>						
<b>Lesion no. 1</b>						
Rate of Repigmentation	0%	1 - 24%	25 - 49%	50 - 74%	75 - 99%	100%

<b>Patient no. 2</b>						
<b>Lesion no. 1</b>						
Rate of Repigmentation	0%	1 - 24%	25 - 49%	50 - 74%	75 - 99%	100%



**APPENDIX C**

**CLINICAL ASSESSMENT OF  
REPIGMENTATION FORM**

Assessment of Repigmentation (CAR)

Please indicate your choice by applying (X);

<b>Patient no. 1</b>						
<b>Lesion no. 1</b>						
Rate of Repigmentation	0%	1 - 24%	25 - 49%	50 - 74%	75 - 99%	100%

<b>Patient no. 2</b>						
<b>Lesion no. 1</b>						
Rate of Repigmentation	0%	1 - 24%	25 - 49%	50 - 74%	75 - 99%	100%

# **CURRICULUM VITAE**



If you don't want this part to be created, you can remove this part.  
If you want, prepare according to the specified format.

2 inches

Page No.

1 inch

1 inch

1.5 inches

## CURRICULUM VITAE

Spacing 24 pt

### NAME

Name Lastname

Spacing 12 pt

### EDUCATIONAL BACKGROUND

Year

Bachelor of Name

Name of Major

Name of Institute

Spacing 12 pt

### WORK EXPERIENCE *(if any)*

Year-present

Position

Institution

Spacing 12 pt

### SCHOLARSHIP *(if any)*

Year

Name of Scholarship

Spacing 12 pt

### PUBLICATION *(if any)*

Write follow APA style pattern. Example:

Abadias, M., Alegre, I., Oliveira, M., Altisent, R., & Viñas, I. (2012). Growth potential of *Escherichia coli* O157:H7 on fresh-cut fruits (melon and pineapple) and vegetables (carrot and escarole) stored under different conditions. *Food Control*, 27(1), 37-44.

\*\*Do not specify the name's title (Mr., Mrs., Ms.)  
If there is a title of the rank, dignity or priest's ranks may or may not be specified.